

**Synthesis and steric structure of 3',3',5'-trimethyl-7',-
'-bis(trifluoromethyl)spiro[1,3,-
-benzodioxaphosphole-2,1'-[2,6,8]trioxa[1λ⁵
]phosphabicyclo[3.2.1]octane] and 3',5'-dimethyl-7',-
'-bis(trifluoromethyl)spiro[1,3,-
-benzodioxaphosphole-2,1'-[2,6,8]trioxa[1λ⁵
]phosphabicyclo[3.2.1]oct[3]ene]**

Mironov V., Baronova T., Mironova E., Dimukhametov M., Krivolapov D., Abdrakhmanova L.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2015 Pleiades Publishing, Ltd. Reactions of hexafluoroacetone with 4-(1,3,-benzodioxaphosphol-2-yloxy)-4-methylpentan-2-one and 4-(1,3,2-benzodioxaphospho-2-yloxy)pent-3-en-2-one afforded cage spiro phosphoranes with a phosphorus-carbon bond, 3',3',5'-trimethyl-7',7'-bis(trifluoromethyl)spiro[1,3,2-benzodioxaphosphole-2,1'-[2,6,8]-trioxa[1λ⁵]phosphabicyclo[3.2.1]octane] and 3',5'-dimethyl-7',7'-bis(trifluoromethyl)spiro[1,3,2-benzodioxaphosphole-2,1'-[2,6,8]trioxa[1λ⁵]phosphabicyclo[3.2.1]oct[3]ene], whose structure was determined by X-ray analysis.

<http://dx.doi.org/10.1134/S1070428015030203>
